

Prepare for Your Doctor's Visit

Get the most out of your appointment.

Benlysta
(belimumab)
Intravenous Use 120 mg/vial
Subcutaneous Use 200 mg/mL

Answer the following questions to create a personalized discussion guide to bring to your next doctor's visit.

My Symptoms

Which of these common lupus symptoms have you experienced? You'll want to tell your doctor about any symptoms you've had, even if they haven't been frequent or recent.

These are not complete lists of all possible symptoms, and BENLYSTA may not help with all of these symptoms. If you're experiencing severe symptoms that require medical attention, please contact your physician immediately.

- | | | |
|---|---|---|
| <input type="checkbox"/> Fever | <input type="checkbox"/> Chest pain | <input type="checkbox"/> Confusion and memory loss |
| <input type="checkbox"/> Fatigue | <input type="checkbox"/> Dry eyes | <input type="checkbox"/> Fingers and toes that turn white or blue when exposed to cold or during stressful periods (Raynaud's phenomenon) |
| <input type="checkbox"/> Joint pain | <input type="checkbox"/> Headaches | <input type="checkbox"/> Other _____ |
| <input type="checkbox"/> Stiffness and swelling | <input type="checkbox"/> Butterfly-shaped rash on the cheeks and bridge of the nose | _____ |
| <input type="checkbox"/> Sun sensitivity | <input type="checkbox"/> Rashes elsewhere on the body | |
| <input type="checkbox"/> Shortness of breath | | |

Have you experienced any of the following signs of lupus nephritis? (optional)

- | | | |
|---|--|---|
| <input type="checkbox"/> Sudden and unexplained swelling, especially in the extremities (feet, ankles, legs, fingers, arms) or the eyes | <input type="checkbox"/> Blood in the urine | <input type="checkbox"/> Increased need to urinate, especially at night |
| | <input type="checkbox"/> Elevated blood pressure | <input type="checkbox"/> Other _____ |
| | <input type="checkbox"/> Foamy or frothy urine | _____ |

How would you describe the impact of these symptoms since your last appointment?

- | | | |
|--|---|--|
| <input type="checkbox"/> One or more of my symptoms has been difficult to manage. | <input type="checkbox"/> One or more of my symptoms has been manageable at times. | <input type="checkbox"/> My symptoms have been manageable. |
| <input type="checkbox"/> I don't know if what I've been experiencing is a symptom. | | |

What is BENLYSTA?

BENLYSTA is a prescription medicine used to treat people 5 years of age and older with active systemic lupus erythematosus (SLE or lupus) or active lupus nephritis (LN) (lupus-related kidney inflammation) who are receiving other lupus medicines. It is not known if BENLYSTA is safe/effective in people with severe active central nervous system lupus.

It is not known if BENLYSTA, given under the skin, is safe/effective in:

- children with SLE under 5 years of age or weighing less than 33 lbs
- children with LN under 18 years of age

It is not known if BENLYSTA, given in a vein (intravenously), is safe and effective in children less than 5 years of age.

Important Safety Information

Do not use BENLYSTA if you are allergic to belimumab or any ingredients in BENLYSTA.

The most important information about BENLYSTA

Immunosuppressive agents, including BENLYSTA, can cause serious side effects. Some of these may cause death.

- **Infections:** fever, chills, pain or burning with urination, urinating often, coughing up mucus, or warm, red, or painful skin or sores on your body. Infections could be serious, leading to hospitalization or death.
- **Allergic (hypersensitivity) reactions:** itching, swelling of the face, lips, mouth, tongue, or throat, trouble breathing, anxiousness, low blood pressure, dizziness or fainting, headache, nausea, or skin rash. Serious allergic reactions can happen the day of, or in days after, receiving BENLYSTA and may cause death.

Please see additional Important Safety Information throughout and accompanying Important Facts About BENLYSTA.

Prepare for Your Doctor's Visit

Get the most out of your appointment.

Benlysta
(belimumab)
Intravenous Use 120 mg/vial
Subcutaneous Use 200 mg/mL

My Treatment Plan

How would you describe your experience with your lupus treatment?

- | | | |
|--|--|---|
| <input type="checkbox"/> I've noticed progress since starting treatment. | <input type="checkbox"/> I've noticed progress but am interested in other options. | <input type="checkbox"/> I'm dissatisfied with my current treatment and am interested in other options. |
| <input type="checkbox"/> I don't know if my treatment is working. | <input type="checkbox"/> I'm not taking any medications for my lupus or lupus nephritis. | |

What are your goals for your treatment plan?

- | | | |
|---|--|--|
| <input type="checkbox"/> I want to better manage my lupus symptoms. | <input type="checkbox"/> I want to reduce severe lupus flares. | <input type="checkbox"/> I want to address high-dose or frequent steroid use. |
| <input type="checkbox"/> I want to better manage my lupus nephritis symptoms. | <input type="checkbox"/> I want to reduce the risk of long-term health concerns. | <input type="checkbox"/> I'm not taking any medications for my lupus or lupus nephritis. |

My Questions About BENLYSTA

Whether you and your doctor have discussed BENLYSTA before, you may still have questions—get the answers you're looking for.

- | | | |
|--|--|--|
| <input type="checkbox"/> How can I tell if BENLYSTA is right for me? | <input type="checkbox"/> What kind of results can I expect from taking BENLYSTA? | <input type="checkbox"/> Which administration option is recommended? |
| <input type="checkbox"/> How long could it take for BENLYSTA to work? | <input type="checkbox"/> What are the potential side effects of BENLYSTA? | <input type="checkbox"/> How often do I take BENLYSTA? |
| <input type="checkbox"/> How would BENLYSTA work with my current treatment plan? | <input type="checkbox"/> How does BENLYSTA work? | <input type="checkbox"/> How was BENLYSTA studied? |

You're all set!

Please print this guide if you'd like a paper copy, or keep it on your phone so you can refer back to it, and discuss your responses with your doctor. To find a doctor near you who can help you manage lupus and learn about BENLYSTA, visit www.benlysta.com/talk-to-your-doctor/find-a-doctor/.

Important Safety Information (cont'd)

The most important information about BENLYSTA (cont'd)

- **Mental health problems and suicide:** thoughts of suicide or dying, attempt to commit suicide, trouble sleeping (insomnia), new or worse anxiety or depression, acting on dangerous impulses, other unusual changes in your behavior or mood, or thoughts of hurting yourself or others.

Before receiving BENLYSTA, discuss with your healthcare provider if you:

- think you have an infection or have infections that keep coming back. Do not use BENLYSTA if you have an infection unless your healthcare provider tells you to.
- have or have had mental health problems such as depression or thoughts of suicide.
- have recently received or may need a vaccination. If you are receiving BENLYSTA, you should not receive live vaccines.
- are taking any medicines, including prescription, over-the-counter, vitamins, and herbal supplements.
- are allergic to other medicines.

Please see additional Important Safety Information throughout and accompanying Important Facts About BENLYSTA.

Prepare for Your Doctor's Visit

Get the most out of your appointment.

Benlysta
(belimumab)
Intravenous Use 120 mg/vial
Subcutaneous Use 200 mg/mL

Important Safety Information (cont'd)

Before receiving BENLYSTA, discuss with your healthcare provider if you: (cont'd)

- are receiving other biologic medicines.
- have or have had any type of cancer.
- have any other medical conditions.
- are pregnant or plan to become pregnant. It is unknown if BENLYSTA will harm your unborn baby. Talk to your healthcare provider about whether to prevent pregnancy while on BENLYSTA. If you choose to prevent pregnancy, you should use an effective method of birth control for at least 4 months after the final dose of BENLYSTA.
- become pregnant while receiving BENLYSTA, talk to your healthcare provider about enrolling in the BENLYSTA Pregnancy Registry. You can enroll in this registry by calling 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/benlysta-belimumab/>.
- are breastfeeding or plan to breastfeed. It is unknown if BENLYSTA passes into your breast milk.

Possible side effects of BENLYSTA

- **Progressive multifocal leukoencephalopathy (PML).** PML is a serious and life-threatening brain infection. PML can result in death or severe disability. Tell your healthcare provider right away if you notice any new or worsening medical problems: memory loss, trouble thinking, dizziness or loss of balance, difficulty talking or walking, or loss of vision.
- **Cancer.** Medicines that affect the immune system, including BENLYSTA, may increase your risk of certain cancers.

The most common side effects of BENLYSTA are nausea, diarrhea, fever, stuffy or runny nose and sore throat, persistent cough, trouble sleeping, leg or arm pain, depression, headache, and pain, redness, itching, or swelling at the site of injection (when given subcutaneously). These are not all the possible side effects of BENLYSTA. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see full Important Safety Information throughout and accompanying Important Facts About BENLYSTA.

 **Benlysta Cares**

Have questions or need assistance?
Call **1-877-4-BENLYSTA**
(1-877-423-6597)

Personalized Support

Benefits & Savings


Exclusive Resources

Whether you need treatment support or financial assistance,
BENLYSTA Cares may be able to help.

Trademarks are owned by or licensed to the GSK group of companies.

©2024 GSK or licensor.
PMUS-BELWCNT240021 July 2024
Produced in USA.

Benlysta  **GSK**



IMPORTANT FACTS ABOUT BENLYSTA

The risk information presented here is not comprehensive. To learn more, talk to your healthcare provider (HCP).

Visit www.BENLYSTA.com or call 1-877-423-6597 to get FDA-approved product labeling, including Medication Guide and Instructions for Use.

What is the most important information I should know about BENLYSTA?

Immunosuppressive agents, including BENLYSTA, can cause serious side effects. Some of these side effects may cause death. Tell your HCP right away if you get any of the following signs or symptoms while receiving BENLYSTA:

- **Infections.** Infections could be serious, leading to hospitalization or death. Symptoms can include: fever, chills, pain or burning with urination, urinating often, coughing up mucus, warm, red, or painful skin or sores on your body.
- **Allergic (hypersensitivity) reactions.** Serious allergic reactions can happen on the day of, or in the days after, receiving BENLYSTA and may cause death. Symptoms can include: itching, swelling of the face, lips, mouth, tongue, or throat, trouble breathing, anxiousness, low blood pressure, dizziness or fainting, headache, nausea, skin rash.
- **Mental health problems and suicide.** Symptoms can include: thoughts of suicide or dying, attempt to commit suicide, trouble sleeping, new or worse anxiety, new or worse depression, acting on dangerous impulses, other unusual changes in your behavior or mood, thoughts of hurting yourself or others.

What is BENLYSTA?

BENLYSTA is a prescription medicine used to treat people 5 years of age and older with active systemic lupus erythematosus (SLE or lupus) or active lupus nephritis (LN) (lupus-related kidney inflammation) who are receiving other lupus medicines. It is not known if BENLYSTA is safe/effective in people with severe active central nervous system lupus.

It is not known if BENLYSTA, given under the skin, is safe/effective in:

- children with SLE under 5 years of age or weighing less than 33 lbs
- children with LN under 18 years of age

It is not known if BENLYSTA, given in a vein (intravenously), is safe and effective in children less than 5 years of age.

Who should not use BENLYSTA?

Do not use BENLYSTA if you are allergic to belimumab or any of the ingredients in BENLYSTA.

What should I tell my HCP before receiving BENLYSTA?

Before you receive BENLYSTA, tell your HCP about all of your medical conditions, including if you:

- think you have an infection or have infections that keep coming back. You should not receive BENLYSTA if you have an infection unless your HCP tells you to. **See “What is the most important information I should know about BENLYSTA?”**
- have or have had mental health problems such as depression or thoughts of suicide.
- have recently received a vaccination or if you think you may need a vaccination. If you are receiving BENLYSTA, you should not receive live vaccines.
- are allergic to other medicines.
- are receiving other biologic medicines or monoclonal antibodies.
- have or have had any type of cancer.
- are pregnant or plan to become pregnant. It is not known if BENLYSTA will harm your unborn baby. You should talk to your HCP about whether to prevent pregnancy while on BENLYSTA. If you choose to prevent pregnancy, you should use an effective method of birth control while receiving BENLYSTA and for at least 4 months after the final dose of BENLYSTA.

- become pregnant while receiving BENLYSTA, talk to your healthcare provider about enrolling in the BENLYSTA Pregnancy Registry. You can enroll in this registry by calling 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/benlysta-belimumab/>.
- are breastfeeding or plan to breastfeed. It is not known if BENLYSTA passes into your breast milk. You and your HCP should talk about whether or not you should receive BENLYSTA and breastfeed.

Tell your HCP about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of BENLYSTA?

BENLYSTA can cause serious side effects including:

- **Progressive multifocal leukoencephalopathy (PML).** PML is a serious and life-threatening brain infection. Your chance of getting PML may be higher if you are treated with medicines that weaken your immune system, including BENLYSTA. PML can result in death or severe disability. Tell your HCP right away if you notice any new or worsening medical problems such as: memory loss, trouble thinking, dizziness or loss of balance, difficulty talking or walking, loss of vision.
- **Cancer.** BENLYSTA may reduce the activity of your immune system. Medicines that affect the immune system may increase your risk of certain cancers.

The most common side effects of BENLYSTA include: nausea, diarrhea, fever, stuffy or runny nose and sore throat, persistent cough, trouble sleeping, leg or arm pain, depression, headache, and pain, redness, itching, or swelling at the site of injection.

These are not all the possible side effects of BENLYSTA. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Keep BENLYSTA and all medicines out of reach of children.

Trademarks are owned by or licensed to the GSK group of companies.



GlaxoSmithKline
Durham, NC 27701

©2024 GSK or licensor.
PMUS-BELBRFS240001 May 2024
Produced in USA.